

GUT-KAVACH

"Gut Health Improvisor"

Newly
Launched

COMPOSITION :

- Probiotic - Bacillus subtilis
- Bacillus coagulans
- Bacillus clausii
 - Prebiotic - Saccharomyces cerevisiae
- Saccharomyces bolardi
- Total viable count : 5×10^9 Cfu per gm (5000 billion spores per kg)
- Gut-Kavach Fortified with -
 - Digested Yeast - (MOS + β -glucan + Yeast cytoplasm + Refined functional carbohydrates)
 - Fructooligosaccharides (FOS) & Inulin
 - Protease
 - Essential oil & UGF

Gut-Kavach is a highly concentrated, well researched symbiotic formulated for the exclusive use in poultry.

ADVANTAGES :

- Improves population of favorable essential microflora in intestine.
- Secretes antibacterial substances.
- Improves production & regeneration of epithelial cells of intestine.
- Reduces issues as 'leaky gut' and improves gut integrity.
- Inactivates bacterial toxins in intestine and reduces ammonia production.
- Increases production of digestive enzymes to improves nutrient digestibility and absorption.
- Improves villi length.
- Improves thickness of mucosal layer forming a better protective barrier of intestinal wall.
- Binds pathogens and reduces their negative impact.
- Helps in promoting the immune system in different ways.
- Secretes micronutrients.

BENEFITS :


- Improves weight gain and FCR
- To control infections and non-specific diarrhea.
- Improves egg production.
- To control dirty egg issues.
- Reduces general mortality
- Improves hatchability and fertility of birds.
- Improves immunity.

INDICATIONS :

- Performance Improvement-- Regular continuous use in feed helps in better performance of birds (Higher Return on Investment)
- Severe Diarrhea-- Controls 75-80% of infectious and noninfectious diarrhea within 5-7 days.
- Dirty Egg Issues-- Helps in solving issues of 'dirty egg'.


DOSAGE:

Gut-Kavach (Water Soluble)	Gut-Kavach (Feed)
Administration through drinking water:	For regular use for improved performance:
Chicks: 20 gm/1000 birds for first 7 days & then 20 gm twice a week Growers/Layers/Breeder/Broilers: 20 gm/1000 birds twice a week	Broiler/Layer/Breeder: 200-250 gm/ton of feed or as directed by nutritionist
Presentation: 1 kg	Presentation: 5 kg



During severe diarrhea:

400-500 gm/ton of feed



During issues of dirty egg:

400-500 gm/ton of feed

